



## Anti-TDP-43 Antibody

**Alternative Names:** TAR DNA-binding protein 43, TARDBP, transactive response DNA binding protein 43 kDa, ALS10

**Catalogue Number:** AB18-10049

**Size:** 100ug

### Background Information

TAR DNA-binding protein 43 (TDP-43) has been shown to bind both DNA and RNA and have multiple functions in transcriptional repression, pre-mRNA splicing and translational regulation. It belongs to the hnRNP protein family and is highly expressed in the pancreas, placenta, lung, genital tract and spleen[1]. Characterisation of transcriptome-wide binding sites revealed that thousands of RNAs are bound by TDP-43 in neurons. TDP-43 regulates alternate splicing of the CFTR gene. The resulting aberrant splicing is associated with pathological features typical of cystic fibrosis.

TDP-43 is predominantly located in the nucleus under normal physiological conditions. However, hyperphosphorylated, fragmented and ubiquitinated forms of TDP-43 have been identified as core components of cytosolic inclusions in sporadic ALS and frontotemporal lobar degeneration (FTLD) [2,3,4,5,6,7,8]. TDP-43 contains a nuclear localising signal (NLS) as well as a nuclear export signal (NES)[8], which enables the shuttling of TDP-43 between the nucleus and the cytosol. Under normal conditions, TDP-43 interacts with mRNAs on which ribosomes are located separately, forming polysomes. Various stresses can induce clustering of ribosomes into a 'stalled' state, resulting in the formation of stress granules (SG) containing TIA-1, G3BP, ataxin-2 and eIF4G1/2.

In the stalled state, transcription is inhibited as a homeostatic response. However, sustained stress and ensuing TDP-43 misfolding creates aberrant SGs and pathogenic TDP-43 aggregates [9]. Moreover, membrane-less organelles in the cytosol formed by the liquid-liquid phase separation of RNPs4 and RNA13 are implicated in TDP-43 proteinopathy. Misfolding and cytosolic mislocalisation also lead directly to a loss of normal TDP-43 function, and the resultant disruption of protein and RNA homeostasis is considered another likely pathogenic mechanism14 in addition to the toxicity of inclusions [9]. Mutations in TDP-43 have been associated with amyotrophic lateral sclerosis, frontotemporal dementia, Parkinson's disease and Alzheimer's disease.

### Product Information

<b>Antibody Type:</b>	Polyclonal	<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG	<b>Species Reactivity:</b>	Human, Mouse, Rat
<b>Immunogen:</b>	Recombinant human TDP-43 N- terminal (aa 1-260)		
<b>Format:</b>	100 µg in 100 µl PBS containing 0.02% sodium azide.		
<b>Storage Conditions:</b>	6 months: 4°C. Long-term storage: -20°C. Avoid multiple freeze and thaw cycles.		
<b>Applications:</b>	IHC, WB, IF, IP, RIP		
	WB 1:500-2000	IHC 1:50-100	IF 1:20-100 IP 1:50-100 RIP 1:20-50



## Additional Information

Subcellular location:	Nucleus	MW: 43kDa
Gene ID	23435	Uniprot ID: Q13148

## References

1. Strong MJ, Volkening K, Hammond R, et al. (2007). TDP43 is a human low molecular weight neurofilament (hNFL) mRNA-binding protein. *Molecular and Cellular Neuroscience*. 35 (2): 320–7.
2. Kwong LK, Neumann M, Sampathu DM, et al. (2007). TDP-43 proteinopathy: The neuropathology underlying major forms of sporadic and familial frontotemporal lobar degeneration and motor neuron disease. *Acta Neuropathologica*. 114 (1): 63–70.
3. Arai, T. et al. TDP-43 is a component of ubiquitin-positive tau-negative inclusions in frontotemporal lobar degeneration and amyotrophic lateral sclerosis. *Biochem. Biophys. Res. Commun.* 351, 602–611 (2006). CAS
4. Neumann M, Sampathu DM, Kwong LK, Truax AC, et al. (2006). Ubiquitinated TDP-43 in Frontotemporal Lobar Degeneration and Amyotrophic Lateral Sclerosis. *Science*. 314 (5796): 130–3.
5. Tan, C. F. et al. TDP-43 immunoreactivity in neuronal inclusions in familial amyotrophic lateral sclerosis with or without SOD1 gene mutation. *Acta Neuropathol.* 113, 535–542 (2007).
6. Igaz, L. M. et al. Enrichment of C-terminal fragments in TAR DNA-binding protein-43 cytoplasmic inclusions in brain but not in spinal cord of frontotemporal lobar degeneration and amyotrophic lateral sclerosis. *Am. J. Pathol.* 173, 182–194 (2008).
7. Tremblay C, St-Amour I, Schneider J, et al. (2011). Accumulation of transactive response DNA binding protein 43 in mild cognitive impairment and Alzheimer disease. *J Neuropathol Exp Neurol.* 70 (9): 788–98.
8. Winton, M. J. et al. Disturbance of nuclear and cytoplasmic TAR DNA-binding protein (TDP-43) induces disease-like redistribution, sequestration, and aggregate formation. *J. Biol. Chem.* 283, 13302–13309 (2008).
9. Yoshitaka Tamaki et al. Elimination of TDP-43 inclusions linked to amyotrophic lateral sclerosis by a misfolding-specific intrabody with dual proteolytic signals. *Scientific Reports: volume 8, Article number: 6030 (2018)*